

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 3 : G01N 33/54, 33/56; C12Q 1/70 C12Q 1/28	A1	(11) International Publication Number: WO 84/ 03564 (43) International Publication Date: 13 September 1984 (13.09.84)
---	-----------	--

(21) International Application Number: PCT/AU84/00039 (22) International Filing Date: 8 March 1984 (08.03.84) (31) Priority Application Number: PF 8348 (32) Priority Date: 8 March 1983 (08.03.83) (33) Priority Country: AU (71) Applicant (for all designated States except US): COMMONWEALTH SERUM LABORATORIES COMMISSION [AU/AU]; 45 Poplar Road, Parkville, VIC 3052 (AU). (72) Inventor; and (75) Inventor/Applicant (for US only) : GEYSEN, Hendrik, Mario [AU/AU]; 10 Janden Close, Knoxfield, VIC 3180 (AU). (74) Agents: SLATTERY, John, Michael et al.; Davies & Collison, 1 Little Collins Street, Melbourne, VIC 3000 (AU).	(81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), DK, FR (European patent), GB (European patent), JP, LU (European patent), NL (European patent), NO, SE (European patent), US. Published <i>With international search report.</i>
--	--

(54) Title: **METHOD OF DETERMINING ANTIGENICALLY ACTIVE AMINO ACID SEQUENCES**

(57) Abstract

A method of detecting or determining a sequence of amino acids which is antigenically active within a known amino acid sequence of a protein or portion thereof comprises the steps of: 1. synthesising a plurality of peptides, each of said peptides comprising a sequence of a plurality of amino acids which corresponds to a sequence within the known amino acid sequence, and the said peptides having overlapping amino acid sequences; 2. contacting each of said peptides with antibody against the protein or portion of interest; and 3. detecting or determining the presence or absence of an antigen-antibody reaction between each of said peptides and said antibody to indicate whether or not said peptide has antigenic activity.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	KR	Republic of Korea
AU	Australia	LI	Liechtenstein
BE	Belgium	LK	Sri Lanka
BG	Bulgaria	LU	Luxembourg
BR	Brazil	MC	Monaco
CF	Central African Republic	MG	Madagascar
CG	Congo	MR	Mauritania
CH	Switzerland	MW	Malawi
CM	Cameroon	NL	Netherlands
DE	Germany, Federal Republic of	NO	Norway
DK	Denmark	RO	Romania
FI	Finland	SD	Sudan
FR	France	SE	Sweden
GA	Gabon	SN	Senegal
GB	United Kingdom	SU	Soviet Union
HU	Hungary	TD	Chad
JP	Japan	TG	Togo
KP	Democratic People's Republic of Korea	US	United States of America

WO 84/03564

PCT/AU84/00039

1

"METHOD OF DETERMINING ANTIGENICALLY ACTIVE AMINO ACID
SEQUENCES"

5 This invention relates to a method of
detecting or determining antigenically active
sequences of amino acids in a protein. The term
"antigenically active" used throughout this
specification is intended to denote any amino acid
sequence which will combine specifically with an
antibody, and includes sequences which also elicit or
10 stimulate the production of antibodies (these latter
sequences are also referred to as being
"immunogenic").

15 As is already well known, an antigen is a
macromolecule such as a protein which is usually
foreign to a human or animal and which is capable of
eliciting the formation of an antibody by the human or
animal, the antibody being a protein synthesized by
the human or animal in response to the presence of the
20 macromolecule. The antibody has specific affinity for
the macromolecule that elicited its synthesis, the
specificity of the antibody being directed against one

25



WO 84/03564

PCT/AU84/00039

2

or more particular sites or amino acid sequences on the macromolecule usually referred to as the "antigenic determinant(s)".

5 It is a primary object of the present invention to detect or determine the sequence or sequences of amino acids which constitute the antigenic determinant(s) of a selected protein, for example, the antigen responsible for eliciting the
10 antibodies which protect against a particular clinical disease such as foot and mouth disease or hepatitis.

According to the present invention there is provided a method of detecting or determining a
15 sequence of amino acids which is antigenically active within a known amino acid sequence of a protein or portion thereof, the method comprising the steps of:

- 20 1. synthesising a plurality of peptides, each of said peptides comprising a sequence of a plurality of amino acids which corresponds to a sequence within the known amino acid sequence, and said peptides having overlapping amino acid sequences;
- 25 2. contacting each of said peptides with antibody against the protein or portion of interest; and
- 30 3. detecting or determining the presence or absence of an antigen-antibody reaction



WO 84/03564

PCT/AU84/00039

3

between each of said peptides and said antibody to indicate whether or not said peptide has antigenic activity.

5

As noted above, the peptides synthesized in accordance with this method have overlapping amino acid sequences, that is, one peptide is related to another peptide in that at least one amino acid is omitted from one end of the sequence of the other peptide and at least one amino acid added to the opposite end of the sequence, with the remaining amino acids being common to both sequences. The amino acids within these overlapping sequences are of course selected so that each sequence corresponds to a sequence within the known amino acid sequence of the protein or portion thereof, as previously described.

10
15

The method of the present invention is based upon the concept that a given antibody specifically recognises a sequence of amino acids which has the particular antigenic activity of interest, and that as a result a particular antigenic determinant can be ascertained by utilising this high degree of specificity of the antibody to identify the specific sequence of amino acids of the particular antigenic determinant from among all the possible combinations thereof.

20
25

30

Several recent publications relate to work aimed at identifying the sequence(s) of amino acids representing the active antigen or antigenic



WO 84/03564.

PCT/AU84/00039

4

determinant of immunogenic proteins. By currently available methods, however, the identification of the active sequence(s) of amino acids within a protein molecule is a long and involved process. The present invention detects the sequences of amino acids which are antigenically active by a screening process using peptides having overlapping sequences as described above.

10 It is believed that an antigenic determinant generally comprises a sequence of about six amino acids in length, and accordingly the sequences of amino acids which are prepared in accordance with the first step of the present invention are preferably
15 sequences of six amino acids. It is to be understood, however, that the present invention is not restricted to sequences of six amino acids although by present indications sequences which are five units in length would seem to be too small to give an easily detected
20 reaction in the subsequent screening procedure. Sequences of nine or more units in length are probably unnecessarily long and thus such sequences are presently not preferred; however, they are encompassed within the broad ambit of the present
25 invention.

In performance of this invention, use is made of previous determination of the amino acid sequence of the protein or protein region believed to
30 carry the antigenic determinant of interest. Amino acid sequences of a large number of proteins are



WO 84/03564

PCT/AU84/00039

5

already known and modern methods of sequencing, for example using recombinant DNA techniques, provide a rapid method of sequencing proteins for which the amino acid sequence is still unknown.

5

10 The importance of this invention lies in its ability to identify the antigenic determinant(s) of a protein as short amino acid sequences with unsurpassed ease. Furthermore, every continuous antigenic
15 determinant of importance can be identified. This information is invaluable in the design of reagents with the necessary selectivity to be used in the diagnosis of clinical diseases in man and animals. A knowledge of the antigenic determinant associated with
20 a particular infectious agent is also essential in the production of peptide vaccines which will confer protection against the disease without many of the deleterious side effects of conventional vaccines. The invention also can be used in the design of very specific therapeutic reagents against receptor sites in the body.

25 One example of the use of the above method in the determination of an antigenic determinant within a protein or portion of a protein is as follows:

- 30 1. Commencing with the known amino acid sequence of the protein or portion of interest, obtained for example from published data, take the order of the first



WO 84/03564

PCT/AU84/00039

6

5 six amino acids (assuming that all
synthesized peptides in this example will be
six amino acids long), starting either from
the amino terminal end or from the carboxy
terminal end, and designate this as
"Sequence one". "Sequence two" is then the
set of six amino acids in order starting at
the second amino acid and proceeding up to
and including the seventh amino acid. By
10 extending this procedure of "dropping" one
amino acid from one end and adding the next
in the known sequence to the other end, the
necessary amino acid sequences for all
15 hexapeptides contained within the protein or
portion of interest are obtained. The total
number of such sequences will be five less
than the actual number of amino acids making
up the protein or portion of interest.

- 20 2. Each sequence determined as described above
is then synthesized. Suitable methods
include the well-known procedures for
peptide synthesis commonly referred to as:
25 (a) solution phase methods; or
(b) solid phase methods; e.g. the
Merrifield technique: Marglin, A. and
Merrifield, R.B., Ann. Rev. Biochem.
39, 841-866 (1970).

30 Preferably, however, the synthesis of



WO 84/03564

PCT/AU84/00039

7

5 the amino acid sequences is effected by a
solid phase method which comprises the use
of a polymeric material such as polyethylene
or polypropylene as the solid-phase carrier,
10 onto which is graft polymerised a vinyl
monomer containing at least one functional
group to produce polymeric chains on the
carrier. The functional groups of these
polymeric chains are then reacted to provide
primary or secondary amino groups of the
15 chains, and these amino groups are then
sequentially reacted with amino acid
residues in appropriate order so as to build
up a desired synthetic peptide. The carrier
is preferably in the form of a solid polymer
rod having a diameter of about 4mm and a
length of about 50mm. A number of such rods
20 can be held in a suitable holder in a 12 x 8
grid whose dimensions correspond to those of
the standard plate used for enzyme-linked
immunosorbent assays (ELISA).

25 3. Depending on the choice of procedure for the
synthesis of each peptide, the peptide is
either already attached to a suitable
support at the completion of the synthesis
procedure, or is coupled to a suitable solid
support in preparation for the assay step.

30 4. The supports carrying the synthesized
peptides are transferred to the wells of a



WO 84/03564

PCT/AU84/00039

8

5 microtitre plate or like apparatus, and each
known sequence is then screened against
known antibodies, using the usual methods to
indicate the presence or absence of an
antigen-antibody reaction in each well.
Examples of these methods include enzyme
linked immunosorbent assay (ELISA) and radio
immuncassay (RIA). Suitable antibodies can
10 either be purchased as commercially
available antisera or prepared in a suitable
host animal in accordance with well known
procedures.

15 Preferably, a computer-based management
program is used to aid in the organisation of the
synthesis of the six-unit long amino acid sequences as
described above. This can be used to achieve a record
of the eventual peptide identity for each location in
each microtitre plate, and in addition allows
20 determination of the order of synthesis to be
undertaken simultaneously on the basis of compatible
conditions of reaction. Finally, such a management
program can be used in conjunction with a program
written for the final evaluation at the antibody
25 screening stage.

30 As will be set out in greater detail in the
Examples, the method outlined in the above steps has
been used to ascertain antigenically active amino acid
sequences of the foot and mouth disease virus (FMDV)
protein VP1. Of course, the present invention can



WO 84/03564

PCT/AU84/00039

9

similarly be applied in determination of antigenic determinants of other virus proteins, such as Hepatitis B virus surface antigen, as well as any other protein for which the sequence is known.

5

The well-known indirect ELISA technique is preferably used for final detection or determination of the presence or absence of an antigen-antibody reaction between the sequences of amino acids and the given antibody. The use of this assay necessitates the preparation of antisera directed against the antibody species to be used in the final testing, for example, anti-human IgG, anti-bovine IgG, and so on. Such preparations are then conjugated with an enzyme, such as horseradish peroxidase, to provide test reagents required for the final testing. Performance of the indirect ELISA assay is carried out by the techniques well known in this field.

20

Where the method of the present invention is performed in a screening test of the type described above, the results obtained with a particular antibody fall into one of the following categories:

25

(a) No positive reaction with any of the prepared amino acid sequences, i.e. complete negative.

(b) Reaction with a single amino acid sequence only. This result represents the ideal case.

30



WO 84/03564

PCT/AU84/00039

10

5 (c) Reaction with a number of amino acid sequences. This observed result can be expected for one of two reasons: one, reaction with an antiserum containing a mixed antibody population, each antibody of which is reacting with a different antigenic site; or two, reaction with amino acid sequences which overlap the immunogenic site. In the first case, further testing will be required to determine the more useful antigenically active amino acid sequences.

10 It will, of course, be appreciated that once the sequence of amino acids which is antigenically active in reacting with a given antibody has been detected or determined, this information can be used
15 in various diagnostic applications, and in the production of vaccines. In fact, where the antigenically active sequence of amino acids corresponding to a particular clinical disease agent has been determined, this can lead to the production
20 of a vaccine which will provide protection against that disease and which comprises one or more synthetically produced sequences of appropriate amino acids which will elicit the desired antibody response.

25 Further features of the present invention are illustrated by way of example only in the following Examples:

30 EXAMPLE 1.

A. Preparation of hexapeptides.



WO 84/03564

PCT/AU84/00039

11

5 The 213-amino acid sequence of VP1 (FMDV, type 0₁) as translated by Kurz, C. et al., Nucleic Acid Research 9, 1919-1931 (1981) was subdivided into all possible hexapeptide units, and each hexapeptide unit was synthesized on a polyethylene support in the same orientation, and with a 2-long amino acid spacer as illustrated in Figure 1.

10 Polyethylene rods immersed in a 5% v/v aqueous solution of acrylic acid were γ -ray irradiated at a dose of 1 Mrad (see Muller-Schulte, D., Horster, F.A., Polymer Bulletin 7, 77-81 (1982)). Using conventional methods of solid-phase peptide chemistry (see Erickson, B.W., Merrifield, R.B. in "The Proteins", 15 Vol.2, 255-257, Academic Press, New York (1976); Meienhofer, J., in "Hormonal Proteins and Peptides", Vol.2, 45-267, Academic Press, New York (1973)), N ^{α} -t-Butyloxycarbonyl-L-Lysine methyl ester was coupled to the polyethylene polyacrylic acid (PPA) via 20 the N -amino group of the side-chain. This was followed by the coupling of Boc-Alanine, to complete a peptide-like spacer. Amino-substitution of the support was determined by reacting NH₂-Lysine(OMe)-PPA with C¹⁴ labelled butyric acid, and was found to be 25 8-10 nmoles/rod.

30 Successive amino acids were added by conventional solid phase peptide synthesis as dictated by the sequence to be synthesized. At the completion of the final coupling reaction, and after removal of the t-butyloxycarbonyl (Boc) protecting group, the



WO 84/03564

PCT/AU84/00039

12

terminal amino group was acetylated with acetic
anhydride in a dimethylformamide (DMF)/triethylamine
mixture. All dicyclohexyl carbodiimide-mediated
coupling reactions were carried out in DMF in the
5 presence of N-hydroxy benzotriazole. The following
side-chain protecting groups were used; O-benzyl for
threonine, serine, aspartic acid, glutamic acid and
tyrosine; carbobenzoxy for lysine; tosyl for
arginine; 4-methyl benzyl for cysteine and
10 1-benzyloxycarbonylamido-2,2,2,-trifluoroethyl for
histidine. Side-chain deprotection was achieved by
treatment with borontris(trifluoroacetate) in
trifluoroacetic acid for 90 minutes at room
temperature (see Pless, J., Bauer, W., Angewante
15 Chemie 85, 142 (1973)). After hydrolysis in
HCl/propionic acid, analysis of sequences included in
the synthesis as controls confirmed that coupling at
each stage had occurred. Before testing by ELISA,
rod-coupled peptides were washed several times in
20 phosphate buffered saline (PBS).

B. Testing of Hexapeptides.

Antigenic profiles for the hexapeptides prepared
25 as described in A. above are shown in Figure 2 as a
vertical line proportional to the ELISA extinction
obtained, over the number giving the location within
the VP1 sequence of the peptide N-terminal amino acid.
Antisera used to produce the different profiles as
30 shown, were as follows:



WO 84/03564.

PCT/AU84/00039

13

- (a) and (b) two different anti-intact virus particle, type O_1 ;
- (c) anti-intact virus particle, as used in (b), after absorption with purified complete virus, type O_1 ;
- 5 (d) anti-virus-subunit, type O_1 ;
- (e) anti-VP1, type O_1 and
- (f) anti-intact virus particle, type C_1 .

10 The enzyme-linked immunosorbent assay was used to test each rod-coupled peptide (RCP) for reactivity with each of the defined antisera described above. RCPs were pre-coated with 10% horse serum, 10% ovalbumin and 1% Tween-80 in PBS, to block
15 non-specific absorption of antibodies, for 1 hour at 37°C. Overnight incubation at 4°C in antiserum diluted 1/40 in the preincubation mixture, was followed by 3 washes in 0.05% Tween-80/PBS. Reaction
20 for 1 hour at 37°C with the appropriate anti-rabbit IgG immunoglobulin coupled to horse radish peroxidase, diluted 1/50,000 in the preincubation mixture, was again followed by extensive washing in PBS/Tween to remove excess conjugate. The presence of antibody was detected by reaction for 45 min with a developing
25 solution (40 mg orthophenylenediamine, 20 μ l of hydrogen peroxide in 100 ml of phosphate buffer, pH 5.0), and the colour produced read in a Titertek Multiscan at 420 nm. After tests, peptides were washed three times at 37°C in 8M urea containing 0.1% 2-mercaptoethanol and 0.1% sodium dodecyl sulphate,
30 followed by several washes in PBS to remove all traces



WO 84/03564

PCT/AU84/00039

14

of bound antibody. The RCPs were then ready for further testing with different antisera.

5 Anti-intact virus particle sera were prepared by immunising rabbits with 50 µg of inactivated, purified virus in complete Freund's adjuvant. The animals were bled 3-4 weeks after the single vaccination. Anti-virus-subunit serum (rabbit) was prepared by immunizing 3 times, 3-4 weeks apart, with 10 µg of
10 acid-disrupted purified virus, initially in complete Freund's and subsequently in incomplete Freund's adjuvant. The polypeptide VP1 was separated from the mixture of proteins obtained from urea disrupted, purified virus, by iso-electric focusing. (see
15 Barteling, S.J., Wagenaar, F., Gielkens, A.L.J., J.Gen.Virol. 62, 357-361 (1982).) After elution from the gel with 8M urea and dialysis against PBS, antiserum was raised as described for 12S above. Antiserum for scan (c) was that used for scan (b), but
20 after absorption with purified virus (1500 µg complete virus was incubated with 1 ml of serum for 72 hours at 4°C), and all virus bound antibodies removed by centrifugation.

25 C. Identification of the virus particle-associated antigenic peptide.

30 Of the four anti-intact virus particle sera tested, scans (a) and (b) show the extremes in the reactivity patterns found. Large quantitative differences in the response to an identical antigen



WO 84/03564

PCT/AU84/00039

15

preparation have been reported before, however, these scans highlight the variability possible in the antibody composition between sera. From an examination of scans (a), (b) and (c), antibody reactive with peptides numbers 146 and 147 are present in whole anti-intact virus sera, but absent after absorption with purified virus. These same antibodies are not observed in the anti-subunit sera, scan (d), and only weakly present in the anti-VP1 sera, scan (e). That some activity was found in the anti-VP1 sera, possibly accounts for the immunizing capacity, albeit weak, of the isolated protein. (see Kleid, D.G., et al., Science 214, 1125-1129 (1981).) It should be noted however that another anti-VP1 serum also tested, while retaining a strong activity in position number 148, showed nothing at positions numbers 146 and 147. The superimposition of scan (c) on scan (b) (absorbed compared to non-absorbed) shows that in addition to the loss of activity to peptides numbers 146 and 147, a reduction in activity to peptides numbers 5, 6 and 206 also occurred. Of these, activity to numbers 5 and 6 was not found in all the anti-intact virus sera tested, whilst number 206 activity was invariably present.

25

From these results, it is concluded that of the sequences found to be reactive, the pair at numbers 146 and 147, that is the hexapeptides Gly-Asp-Leu-Gln-Val-Leu (G - D - L - Q - V - L) and Asp-Leu-Gln-Val-Leu-Ala (D - L - Q - V - L - A), constitute the principal loci, with a lesser

30



WO 84/03564.

PCT/AU84/00039

16

contribution from the locus at number 206, consistent with the observations of others. However, with respect to the loci at numbers 146-7, we do not distinguish between the two possibilities; one, that the active element is five amino acids long, i.e. the sequence common to both Asp-Leu-Gln-Val-Leu (D - L - Q - V - L); or two, that the active element is seven amino acids long, i.e. the combination of the two hexapeptides Gly-Asp-Leu-Gln-Val-Leu-Ala (G - D - L - Q - V - L - A).

EXAMPLE 2

A. Preparation of hexapeptides.

The 212 amino acid sequence of VP1 (FMDV type A₁₀ or A₆₁) as given by Bachrach, H.L., et al, Office International des Epizootics was subdivided into 207 hexapeptides. These hexapeptides were synthesised as described in Example 1 above with the exception that the side chain of arginine was protected by the p-methoxybenzene sulphonyl group.

B. Testing of hexapeptides.

Antigenic profiles for the hexapeptides are shown in Figure 3. The antisera used to produce the profiles were:-

- (a) anti-intact virus particle type A₁₀
- (b) anti-intact virus particle as used in (a) after adsorption with purified complete FMDV type A₁₀.



WO 84/03564

PCT/AU84/00039

17

The testing of the hexapeptide and preparation of the sera were essentially as described in Example 1.

5 C. Identification of virus-particle associated active element.

By reasoning identical to that used in Example 1 it is concluded that the hexapeptides Gly-Asp-Leu-Gly-Ser-Ile (G - D - L - G - S - I) and
10 Asp-Leu-Gly-Ser-Ile-Ala (D - L - G - S - I - A) are the principal loci for the antigenic determinant of the A-type of FMDV.

As in the case of FMD virus, type O₁,
15 described in Example 1, we did not distinguish between these two sequences, and accordingly it is concluded that it is possible that the active sequence is five amino acids long, i.e. Asp-Leu-Gly-Ser-Ile (D - L - G - S - I), or that it is seven amino acids long, i.e.
20 Gly-Asp-Leu-Gly-Ser-Ile-Ala (G - D - L - G - S - I - A).

EXAMPLE 3.

25 A. Preparation of hexapeptides.

The 210 amino acid sequence of VP1 (FMDV type C₁) as given by Robertson, H.L., et al., Journal of Virology, 46, 311-316 (1983) was subdivided into 205
30



WO 84/03564

PCT/AU84/00039

18

hexapeptides. These hexapeptides were synthesised as described in Example 2 above.

B. Testing of hexapeptides.

5 Antigenic profiles for the hexapeptides are shown in Figure 4. The antisera used to produce the profiles were:-

- (a) anti-intact virus particle type C₁
- (b) anti-intact virus particle as used in (a)
- 10 after absorption with purified complete FMDV type C₁.

The testing of the hexapeptide and preparation of the sera were essentially as described in Example 1.

15 C. Identification of virus-particle associated active element.

By reasoning identical to that used in Example 1 it is concluded that the hexapeptide Asp-Leu-Ala-His-Leu-Thr (D - L - A - H - L - T) is the principal locus
20 for the antigenic determinant of the C-type of FMDV.

These results clearly show the potential of a systematic scanning of a polypeptide sequence. They
25 point out the likely location of the active determinant encompassed within the peptide with which Bittle, J.L. et al, Nature 298, 30-33 (1982), obtained the successful protection in guinea pigs to a subsequent challenge by FMDV.

30



WO 84/03564

PCT/AU84/00039

19

It will, of course, be recognised that many variations and modifications may be made to the detailed description of the method of the present invention given above without departing from the
5 method of the invention as broadly described herein.



WO 84/03564

PCT/AU84/00039

20

CLAIMS:

1. A method of detecting or determining a sequence of amino acids which is antigenically active within a known amino acid sequence of a protein or portion thereof, the method comprising the steps of:
 1. synthesising a plurality of peptides, each of said peptides comprising a sequence of a plurality of amino acids which corresponds to a sequence within the known amino acid sequence, and the said peptides having overlapping amino acid sequences;
 2. contacting each of said peptides with antibody against the protein or portion of interest; and
 3. detecting or determining the presence or absence of an antigen-antibody reaction between each of said peptides and said antibody to indicate whether or not said peptide has antigenic activity.
2. A method according to claim 1, wherein said plurality of peptides overlap by one amino acid.
3. A method according to claim 1 or claim 2, wherein each of said plurality of peptides is a sequence of from five to nine amino acids.



WO 84/03564

PCT/AU84/00039

21

4. A method according to claim 3, wherein each of said plurality of peptides is a sequence of six amino acids.

5. A method according to claim 1 wherein each of said plurality of peptides is synthesised on a solid phase carrier, and subsequent steps of the method are carried out with each peptide remaining coupled to said solid-phase carrier.

6. A method according to claim 5, wherein said solid phase carrier is in the form of a solid polymer pin or rod.

7. A method according to claim 5 or claim 6, wherein said solid phase carrier is a polymeric material selected from the group consisting of polyethylene or polypropylene, having a vinyl monomer containing at least one functional group graft polymerised thereto to produce polymeric side-chains on the carrier.



WO 84/03564

PCT/AU84/00039

1/5







FIG 1

WO 84/03564

PCT/AU84/00039




2/5



WO 84/03564

PCT/AU84/00039

3/5



WO 84/03564

PCT/AU84/00039

4/5




FIG 3a




FIG 3b



WO 84/03564

PCT/AU84/00039

5/5




FIG 4a




FIG 4b



INTERNATIONAL SEARCH REPORT

International Application No PCT/AU 84/00039

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ³		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int. Cl. ³ G01N 33/54, 33/56, C12Q 1/70, 1/28		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
IPC	G01N 33/54, 33/56, C12Q 1/70, 1/28	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁵		
AU: IPC as above		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁶	Citation of Document, ¹⁵ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
Y	EP, A, 0044710 (SCRIPPS CLINIC AND RESEARCH FOUNDATION) 27 January 1982 (27.01.82)	
Y	Scientific American, February 1983, R.A. Learner, "Synthetic Vaccines", see pages 48 to 56	
Y	Pro. Natl. Acad. Sci. USA, Vol. 77, No. 9, issued September 1980, Walter G. et al, "Antibodies specific for the carboxy-andamino-terminal regions of simian virus 40 large tumor antigen", see pages 5197 to 5200	
Y	AU, A, 12553/83 (BIOGEN N.V.) 29 September 1983 (29.09.83)	
Y	Annual Review: BIOCHEMISTRY, Vol. 39, issued 1970, Marglin A. and Merrifield R.B., "Chemical Synthesis of Peptides and proteins", see pages 841 to 866	
<p>* Special categories of cited documents: ¹⁶</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ¹⁹	Date of Mailing of this International Search Report ²⁰	
30 May 1984 (30.05.84)	(05-06-84) 5, JUNE 1984	
International Searching Authority ¹	Signature of Authorized Officer ²⁰	
Australian Patent Office	A.S. Moore <i>A.S. Moore</i>	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON
INTERNATIONAL APPLICATION NO. PCT/AU 84/00039

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Members			
EP	44710	AU 72836/81	DK 3184/81	EP	44710
		ES 504033	FI 812092	IL	63224
		JP 57118545	NO 812341	PT	73346
		ZA 814637			
AU	12553/83	DK 1390/83	EP 90581	IL	68201
		JP 58213723	NO 831098	PT	76451
		ZA 831854			

END OF ANNEX